

Extravasation Treatment Decision Table



Medication Name	% Swelling and Infusate Component	Action
<p>Red list <u>Vasoactive</u> medications (Dopamine, Dobutamine, Epinephrine, Norepinephrine, Vasopressin, Terlipressin and Phenylephrine)</p>	<p>Extravasation of any % of swelling</p>	<p>IMMEDIATE consult to the Vascular Access Team and Provider to determine treatment plan and use of Phentolamine.</p>
<p>Red list Infusate *(All except vasoactive medications)</p>	<p>Extravasation \geq 30%</p>	<p>Treat with Hyaluronidase per provider order.</p>
	<p>Extravasation $<$ 30%</p>	<p>Clinical evaluation of the extravasation site by the Vascular Access Team and provider to determine if Hyaluronidase is clinically indicated. Decision criteria include imminent skin loss, and / or peripheral circulation impairment (compartment syndrome).</p>
<p>Yellow or Green list Infusate</p>	<p>Extravasation \geq 30%</p>	<p>Clinical evaluation of the extravasation site by the Vascular Access Team and provider, to determine if Hyaluronidase is clinically indicated. Hyaluronidase treatment usually NOT indicated.</p>
	<p>Extravasation $<$ 30%</p>	<p>No treatment indicated.</p>